

READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON HORSES OR DOGS
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Hazard: If absorbed through skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. All handlers must wear long sleeved shirt and long pants, socks and shoes. Remove and wash contaminated clothing before reuse.

For external use only. Do not use on animals under three months of age or on sick, ill, debilitated or medicated pets. Do not apply to horses or dogs intended for slaughter. Do not use on Cats. Sensitization may occur after using REPPEL-X product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

FIRST AID	
IF ON SKIN	• Take off contaminated clothing.
IF ON CLOTHING	• Rinse skin immediately with plenty of water for 15-20 minutes.
	• Call a poison control center or doctor for treatment advice.

See the product container or label with you after calling a poison control center or doctor, or going for treatment. In case of medical emergency, call 1-800-234-2299 24 hours a day, 7 days a week.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into streams, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that water or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees exposed to direct treatment or drifting spray or wash. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

KILLS AND REPELS: Stable flies, horse flies, face flies, deer flies, house flies, horn flies, mosquitoes, gnats, ticks, fleas, chiggers, and lice.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

Remove animals prior to treatment. Wash treated surfaces to dry before allowing animals back into quarters treated with this product. Do not apply directly to any animal destined for riding or slaughter.

Do not use on horses older than 2 months of age.

Do not contaminate water, feed or healthful, wild or milking animals.

For use on a surface spray:

- Do not apply as a broadcast treatment to below surface of walkable areas such as manure pits, dogrun, corral, schools, hospital and morning rooms.
- Do not apply this product in a way that will contact animals, children, or pets either directly or through drift. Remove pets, kids, and cover hot aquifers before use.
- Do not allow adults, children, or pets to enter until spray has dried.

Manufactured by **FARNAM COMPANY, INC.**
 5201 East Swallowford Road, 2008 • Schuylburg, IL 62157
 www.farnam.com
 If case of emergency or for product use information, call 800-234-2299

Ready To Use



REPEL-X

Insecticide & Repellent

**Kills and Repels: Six Fly Species,
 Ticks, Gnats and Mosquitoes**

For Use on Horses, Dogs and Their Premises



ACTIVE INGREDIENTS:	
Pyrethrin	0.00%
Piperonyl Butoxide*	0.50%
Permethrin**	0.10%
OTHER INGREDIENTS:	99.30%
TOTAL:	100.00%

* (Insecticide) 6-propylpiperonyl ether and related compounds
 ** (2-phenoxyphenyl) methyl 3-(4-chlorophenyl) 2,2-dimethylpropanoate
 (insecticide, desferrioxalate, mch, 20%) (4-1-05 and 100-015) (1-1-05)

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for additional Precautionary Statements

Net Contents 1 Qt (.946 L)

FOR HORSES USE FULL STRENGTH: This one-step insecticide repellent may be applied with a trigger spray applicator, or as a wipe, fill and repeat stable fly, horse fly, face fly, deer fly, house fly, horn fly, mosquito, gnats, ticks, fleas, chiggers, and lice. Apply at a maximum rate of 1 ounce per hundred pounds of weight. Do not spray to the point of run-off.

DIRECTIONS FOR HORSES SPRAY USE: Remove excess dirt and dust. Apply light spray mist to coat while brushing lightly against fly of the body. Avoid spray in eyes and mucous membranes. Apply with sprayer or diffuser as directed.

DIRECTIONS FOR WIPES-ON USE: Thoroughly brush horse to remove excess dirt and dust. Extensively dry horse without abandonment, bleed and allowed to dry before applying spray. Use a sponge, clean wet cloth, or mitt. Apply liberally over areas to be protected. Pay special attention to legs, belly, sheath, neck, and facial areas. Avoid eyes and mucous membranes.

TO KILL AND REPEL FLYING INSECTS IN STABLES, QUARTERS: Use this product in horse and cattle feed and dairy barns, milk houses, poultry houses, swine quarters, kennels, schoolyards, pens and shelters to kill and repel stable fly, horse fly, face fly, deer fly, house fly, horn fly, mosquito, gnats, ticks (including deer ticks), fleas, chiggers, and lice. Apply undiluted as a surface spray. Do Not Spray! Use a great sprayer adjusted to deliver a coarse wet spray. Apply as a coarse wet spray using 1 pint per 1,000 square feet. Direct the spray whenever flies congregate or drift, with surfaces are wetted. Treat around drainage, feed storage rooms, alleysways and windows.

Repeat treatment daily or at intervals needed to give adequate protection.

FLIES AND TICKS ON DOGS: Start spraying at the animal's head, working the eyes and mouth in your spray. Make sure the animal's entire body is covered including the legs, underbelly and tail. While spraying, lift the hair so that the spray will penetrate to the skin. Make sure spray with thoroughly. Avoid contact with genitalia. Spray at the rate of 1 or 2 measures of the trigger sprayer per pound of body weight. For longer spray spray will form, not saturated. Do not allow animals to become saturated. If saturated, bathe with water once per day. Do not use on puppies under three months of age. Do not use on cats. Sensitization may occur after using any pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

PET AREAS: Not to be used for broadcast treatment. To control flies and prevent dog ticks on premises as a spot treatment, directly spray infested areas, pet beds, feeding quarters, kennel crates and crates, along and behind beds/benches, readings, window and door frames, and coat areas of floor and floor covering. Pet bedding should be placed in animal's quarters for regular treatment. Document treatment of animals is recommended. Reapply every day.

STORAGE AND DISPOSAL

Do not contaminate water, feed, or feed to storage or disposal. **PESTICIDE STORAGE:** Store in a cool, dry (preferable locked) area that is inaccessible to children and pets and away from food, aquatic, or open flames. Store pesticides away from food, pet food, feed, seed, fertilizer, and veterinary supplies. **PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Non-refillable containers: Do not reuse or refill the container. If empty, place in trash or other for recycling if available. If partly filled Call your local solid waste agency for disposal instructions. Never place chemical product down any drain or into any stream.



Your Partner in Horse Care™

EPH Reg. No. 270-204

EPH Reg. No. 2704-75-1

Product Code: 10330

RNA: 300122973 • 17-11983

©2006, 2010, 2016, 2017 Farnam

Companies, Inc.

Repeat X, Farnam with design
 and Your Partner in Horse Care are trademarks of
 Farnam Companies, Inc.

